

Fakultät für Mathematik, Institut für Mathematische Stochastik
 Prof. G. Christoph, Lora Todorova

Tutorial Stochastic Processes, (Serie 4)

Problems with * are additional problems.

- 16.) Suppose $\Omega = \{1, 2, 3, 4, 5, 6\}$ with $P(\{i\}) = 1/6$ for $i = 1, 2, 3, 4, 5, 6$.

Defin $X(\omega) = \omega$ = face value on die with $P(X = i) = 1/6$ for $i = 1, 2, 3, 4, 5, 6$

$$\text{and } Y(\omega) = \begin{cases} 2 & \text{if } \omega \in \{1, 3, 6\} \\ 3 & \text{if } \omega \in \{2, 5\} \\ 5 & \text{if } \omega = 4 \end{cases} . \quad \text{Find } Z = E(X|Y) !$$

- 17.) a) Let X be uniform on $(0, 1]$, $C_1 = (0, 1/2]$, $C_2 = (1/2, 1]$,

and $\mathcal{F}_2 = \sigma(C_1, C_2)$. Find $V = E(X|\mathcal{F}_2)$!

b) Remember Example 3.2 with $\mathcal{F}_4 = \sigma(A_1, A_2, A_3, A_4)$.

Here $C_1 = A_1 \cup A_2$, $C_2 = A_3 \cup A_4$

$$\text{and } Z(\omega) = E(X|\mathcal{F}_4) = \begin{cases} 1/8 & \text{if } \omega \in A_1 \\ 3/8 & \text{if } \omega \in A_2 \\ 5/8 & \text{if } \omega \in A_3 \\ 7/8 & \text{if } \omega \in A_4 \end{cases} . \quad \text{Find } U = E(Z|\mathcal{F}_2) !$$

- 18.) Let $(W(t), t \geq 0)$ be a standard Wiener Process and $(N(t), t \geq 0)$ a homogenuos Poisson Process mit dem Parameter $\lambda > 0$. Prove that

$$(a) W^2(t) + 2W(t) - t,$$

$$(d) N(t) - \lambda t,$$

$$(b) W^3(t) - 3tW(t),$$

$$\text{and (e*) } (N(t) - \lambda t)^2 - \lambda t,$$

$$(c^*) \exp\{-\theta W(t) - \frac{1}{2}\theta^2 t\}$$

$$(f^*) \exp\{-\theta N(t) + \lambda t(1 - e^{-\theta})\}$$

are martingales with respect to its natural filtration

$$\mathcal{F}_t = \mathcal{F}(W(u), 0 \leq u \leq t) \text{ or } \mathcal{F}_t = \mathcal{F}(N(u), 0 \leq u \leq t) .$$

(g) Show that $W^3(t)$ and $N(t)$ do not be martingales.

- 19.) Let $(W(t), t \geq 0)$ be a standard Wiener Process. Calculate

$$(4.1) E \int_0^t W_s dW_s = 0 \text{ and}$$

$$(4.2) E(\int_0^T W_s dW_s)^2 = \int_0^T E(W_s^2) ds$$

- 20*.) Try to find a stochastic process A_t such that $(W_t^4 + A_t, \mathcal{F}_t)$ is a martingale with respect to the filtration $\mathcal{F}_s = \sigma(W_u, 0 \leq u \leq s)$, $s \geq 0$.

Hint: First calculate $E((W_t - W_s) + W_s)^4 | \mathcal{F}_s)$ for $s < t$.